

Relativistic Whistle: High Order Harmonics Induced by the Ultra-Intense Laser Pulse Propagating inside the Fiber

S. V. Bulanov,^{1,*} T. Zh. Esirkepov,^{2,†} N. M. Naumova,¹ and I. V. Sokolov³

¹*General Physics Institute RAS, Vavilov str. 38, Moscow 119991, Russia*

²*Moscow Institute of Physics and Technology, Institutskij per. 9, Dolgoprudny Moscow region 141700, Russia*

³*University of Michigan, Ann Arbor, MI 48109-2143, USA*

(Dated: May 22, 2002)

A propagation of an ultra-intense short laser pulse in a fiber is investigated with two dimensional Particle-in-Cell simulations. The fiber is a narrow hollow channel with walls consisting of overdense plasma. In the nonlinear interaction of the laser pulse with fiber walls high order harmonics are generated. Sufficiently high harmonics, for which the fiber walls are transparent, propagate outwards at certain angle. This is a scheme of a generator of ultra-short pulses of coherent light with a very short wavelength.

PACS numbers: 42.65.Ky, 52.25.Os, 52.65.Rr, 52.27.Ny

Keywords: high order harmonics, short laser pulse, fiber, Particle-in-Cell simulation

I. INTRODUCTION

Generation of high order harmonics of the electromagnetic radiation during an interaction of an ultra-intense laser pulse with underdense and overdense plasmas is a manifestation of one of the most basic nonlinear processes in physics. High order optical harmonics have been observed in the laser plasma interaction for the laser intensity ranging from a moderate level to relativistic intensities.

High order harmonics attract a great attention due to a wide range of their applications for the diagnostics, the UV and coherent X-ray sources, lithography etc. (see Refs. [1, 2, 3, 4]). Recently, the high order harmonics polarization properties have been used in Ref. [5] to detect the strongest magnetic field generated in the laser plasmas.

Different physical mechanisms of high order harmonics generation have much in common because of a property of nonlinear systems to react in an anharmonic manner to a periodic driving force. On the other hand, specific realization of this property depends on the circumstances of the laser-matter interaction and mainly on the laser intensity. At moderate intensities, generation of high order harmonics is caused by an atom anharmonic response to an oscillating electric field. At relativistic intensities, the electron quiver energy becomes higher than the rest mass energy and the Relativistic Nonlinear Optics comes into play [4]. In this regime high order harmonics generation is due to a nonlinear dependence of the particle mass on its momentum and due to electron density modulations in the electromagnetic wave. In the present paper we discuss high harmonic generation in collisionless plasmas by a relativistically strong electromagnetic wave.

In the underdense plasmas an ultra-intense laser pulse causes high harmonics generation through a parametric excitation of electromagnetic and electrostatic waves with different frequencies. In an interaction with overdense plasmas the laser radiation (partially) reflects back at the plasma-vacuum interface, in the case of sharp plasma boundary, or at a critical density surface, in the case of gradual density profile. The reflecting layer in the plasma is dragged by the electromagnetic wave back and forth in the direction of the incident light, as well as in the transverse direction, forming an oscillating mirror (see Refs. [6, 7, 8, 9, 10, 11, 12]). The light reflected by the oscillating mirror contains odd and even harmonics, whose polarization and amplitudes depend on the incidence angle, intensity and a polarization of the laser pulse.

High efficiency of the laser energy transformation into the energy of high harmonics can be achieved in a scheme, in which the laser pulse propagates inside a hollow channel [13]. In this case, if the channel is sufficiently long, the laser pulse undergoes a multiple reflection at the plasma-vacuum interface, and a portion of the laser pulse energy transforms into the energy of high harmonics at each reflection event. If the laser pulse propagates inside a hollow channel in a tubiform fiber, the radiation with sufficiently high frequency propagates through the fiber walls. The

*Also at Advanced Photon Research Center, JAERI, Kizu minami, Kyoto-fu 619-0215, Japan; Electronic address: bulanov@plasma.fpl.gpi.ru

†Electronic address: timur@nonlin.mipt.ru

fiber walls are transparent for harmonics, whose indices are greater than some critical value. Only these harmonics are emitted from the fiber at certain angle with respect to the fiber axis. This approach allows one to design a high frequency radiation source with controlled properties, which depend on the fiber diameter and the fiber wall thickness. We call this source “the relativistic whistle”.

In this paper we investigate a propagation of an ultra-intense short laser pulse in a fiber, with an aim to study the properties of the high harmonic radiation. Here we use the two dimensional approximation, leaving the investigation of the 3D problem for forthcoming publications.

II. MATCHING CONDITIONS

In the two-dimensional case the fiber corresponds to two thin foils of a finite length. The foils are parallel to each other, and to the x -axis in the x, y -plane. Let the wall thickness of the foil is l , and a distance between the foils equal to L . We consider the foils to be made of the collisionless overdense plasma with the electron density n_e . Such a configuration can be formed by an ultra-intense femtosecond laser pulse interacting with the solid density fiber, when the fiber wall material is ionized during the first half-period of the laser pulse, and the length of the laser pulse is much shorter than the timescale of a hydrodynamic expansion of the plasma. In the framework of this approximation we can assume the ions to be at rest.

From the dispersion equation, $\omega^2 = k^2 c^2$, where ω is the wave frequency and $k^2 = k_x^2 + k_y^2$, we find the group velocity of the wave inside the channel, $v_{gr} = \partial\omega/\partial k_x = (c/\omega)(\omega^2 - \pi^2/L^2)^{1/2} = c \cos \theta$, where $\tan \theta = k_y/k_x = (4L^2/\lambda^2 - 1)^{-1/2}$ and $\lambda = 2\pi c/\omega$. The matching conditions for the wave inside the channel, and the wave radiated outwards, show that all the outgoing waves propagate at the angle θ with respect to the channel axis, independently of the wave frequency.

The fact that all the outgoing waves have the same angle of propagation can be demonstrated in a different way. After the Lorentz transform into the reference frame moving at a speed of v_{gr} along the x -axis, the x -component of the fundamental mode wave vector vanishes: $k'_x = 0$ (see for details Refs. [6, 9, 14]). The problem becomes one-dimensional. Therefore high frequency waves excited by the fundamental mode at the plasma-vacuum interface also have zero x -component of the wave vector in this reference frame. They propagate outwards in the direction perpendicular to the fiber walls. Performing the inverse transform back to the laboratory reference frame, we obtain that all the outgoing waves propagate at the same angle θ with respect to the channel axis, independently of the wave frequency. Their energy is localized inside a slab co-moving with the laser pulse along the x -axis. The slab length is equal to the length of the laser pulse. Its transverse size depends on the effective length of the interaction. According to the selection rules of the harmonic generation at each wall (see Refs. [6, 8, 9]), the s -polarized fundamental mode in the fiber generates s -polarized odd harmonics and p -polarized even harmonics of radiation. The p -polarized fundamental mode generates only p -polarized harmonics of radiation.

The amplitude of the n -th harmonic can be found in the approximation of an infinitely thin walls of the fibre (see Ref. [9]). It is $E^{(n)} = K^{(0,n)} E^{(0)}$, where $E^{(0)}$ and $E^{(n)}$ are amplitudes of the fundamental mode and of the n -th harmonic, respectively. The transformation coefficient $K^{(0,n)}$ is given by $K^{(0,n)} \approx [(a_0/\epsilon) \sin \theta]^n$, in the limit $n\omega/\omega_{pe} \ll \epsilon$. Here $a_0 = eE^{(0)}/m_e\omega_0 c$ is the dimensionless amplitude of the fundamental mode, $\epsilon = 2\pi n_0 e^2 l/m_e\omega_0 c$ is the dimensionless parameter (relativistic transparency measure), ω_0 is the frequency of the fundamental mode.

III. PARAMETERS OF THE COMPUTER SIMULATIONS

We perform computer simulations using the two-dimensional version of Particle-in-Cell relativistic electromagnetic code REMP, based on ‘density decomposition’ scheme [15]. Computation box has a size $45\lambda \times 20\lambda$. Since the problem of high harmonic generation requires substantially high spatial resolution, the computation mesh has 50 cells per λ . The number of quasiparticles per cell is equal to 16 in the plasma region. The channel walls have a length $l_{ch} = 25\lambda$ and width 2λ , a distance between the walls is $L = 0.7\lambda$. The ions are assumed to be immobile and the electron density is equal to $8n_{cr}$, where the critical density is $n_{cr} = \omega_0^2 m_e / 4\pi e^2$. Since the channel diameter is less than the laser light wavelength, only the s -polarized electromagnetic wave (whose electric field is directed along the z -axis) can propagate inside the channel. For the parameters chosen, only harmonic, whose index is higher than 4, can propagate through the walls. This conclusion follows from the boundary conditions for the electromagnetic wave, which is obliquely incident on the plasma-vacuum interface, at the angle θ . The s -polarized laser pulse is initialized in a vacuum region at the left hand side from the channel entry. The pulse aperture is $f/1$, its length is equal to 7.5λ , and its dimensionless amplitude at the 1λ focus spot is $a_0 = 3$, which corresponds to the intensity $I \approx 10^{19} \text{W/cm}^2$. These parameters are close to the parameters of the laser in the CUOS at the University of Michigan [4]. We notice here that the pulse focusing into the 1λ spot is discussed in Ref. [4]. In our simulation model the entry of the channel is smoothed to




FIG. 1: The electromagnetic energy density and the plasma density distribution at $t = 15 \times 2\pi/\omega_0$ when the pulse enters into the fiber.




FIG. 2: The electromagnetic energy density distribution in the x, y plane at $t = 20 \times 2\pi/\omega$ in frame (a), $t = 35 \times 2\pi/\omega$ in frame (b), and at $t = 50 \times 2\pi/\omega$ in frame (c).

decrease the laser pulse reflection. Fig. 1 shows the focusing laser pulse and the fiber. We see that the smoothing of the channel entry provides almost reflectionless matching of the laser pulse and the channel.

IV. RESULTS OF PIC SIMULATIONS

When the laser pulse propagates inside the channel a portion of its energy is reflected back due to the process similar to the stimulated Raman scattering in the plasma of the channel walls; a portion of its energy is transformed into the energy of high harmonic waves radiated outwards; the remaining part of the laser pulse propagates through the channel toward the end of the fiber, then it goes out as a diverging electromagnetic wave. These successive phases are seen in Fig. 2, where we present the electromagnetic energy density distribution in the x, y -plane at different times.

In Fig. 3 we show the distribution of the z -component of the electric field in the x, y -plane. We see the laser pulse




FIG. 3: The distribution of the z -component of the electric field in the x, y -plane at $t = 45 \times 2\pi/\omega_0$.




FIG. 4: Spectrum of the s -polarized harmonics. The dashed line denotes the dependency $k_y \approx 0.8k_x$. Each peak corresponds to a harmonic with index $n = (k_x^2 + k_y^2)^{1/2}$.

inside the channel and the high harmonics aside the channel. In this figure all the harmonics have odd indices and they are s -polarized, as noticed above. The spatial region filled by high harmonics has a length equal to the laser pulse length and the transverse size (along the y -axis) equal to $\delta t c \sin \theta$, where $\sin \theta \approx 1/2$ and $\delta t \approx 20 \times 2\pi/\omega_0$.

We perform the spatial Fourier transform of the z -component of the electric field inside a sub-domain ($18\lambda < x < 28\lambda$; $3\lambda < y < 10\lambda$) at $t = 30 \times 2\pi/\omega_0$. The resulting spectrum of the s -polarized light inside this sub-domain is presented in Fig. 4. We see the odd harmonics with the harmonic index $n = (k_x^2 + k_y^2)^{1/2} \geq 5$. The spectrum peaks are arranged along the straight line given by $k_y \approx 0.8k_x$, i. e. the angle of propagation is smaller than $\arccos(1 - \pi^2 c^2 / \omega_0^2 L^2)^{1/2}$. It means that the effective channel width is larger than 0.7λ . The increase of the channel width is apparently due to the action of the ponderomotive force on the electrons at the plasma-vacuum interface.

From the expression $m_e c^2 \nabla(1 + a^2)^{1/2} \approx 4\pi n_0 e^2 \delta y$, which corresponds to the balance between the ponderomotive force and the force due to electric charge separation, we find that the channel width becomes larger by $2\delta y \approx 0.1\lambda$. Assuming now the channel width to be equal to 0.8λ we obtain the propagation angle $\theta_{eff} \approx \arctan(0.8) \approx 38.7^\circ$, in agreement with the dependence seen in Fig. 4.

In Fig. 5 we present the distribution of the z -component of the magnetic field at the same point in time as Fig. 3. We see the high harmonics with even indices. They are p -polarized. As noticed above, the fundamental mode of the laser pulse inside the channel does not contain the p -polarized component and we see only the high harmonics inside and aside the channel, as well as the low frequency surface mode in the vicinity of the outer plasma-vacuum interfaces on both sides of the fiber. The even-index high harmonics outside the channel are localized in the same region as the odd-index harmonics.

The spectrum of the even-index high harmonics calculated for the z -component of the magnetic field in the same sub-domain and at the same point in time as in the case of the odd-index harmonics is presented in Fig. 6. We see the even harmonics with the harmonic index $n \geq 6$. As in the previous figure, the spectrum peaks are arranged along the same line $k_y \approx 0.8k_x$.




FIG. 5: The distribution of the z -component of the magnetic field in the x, y -plane at $t = 45 \times 2\pi/\omega_0$.




FIG. 6: Spectrum of the p -polarized harmonics.

V. CONCLUSION

The high harmonic generation in the interaction of the ultra-intense short laser pulse with a narrow fiber is demonstrated with PIC simulation. Due to nonlinear interaction of the laser pulse with the fiber walls, the quivering back and forth electrons form the oscillating mirrors. The high order harmonics are generated as a result of a multiple reflection of the laser light from the oscillating mirrors. The fiber walls are opaque for the electromagnetic wave with a frequency below some critical value, which is defined by the electron density of the fiber walls and the channel width. As a result, only sufficiently high harmonics, for which the fiber walls are transparent, propagate outwards at certain angle, independently of the harmonic index. In addition, we observe the excitation of the p -polarized low frequency mode, which is localized at the outer surface of the fiber. The longitudinal scale of the low frequency mode is of the order of the laser pulse length. It provides an example of the nonlinear rectification of the light. The high harmonics energy is localized in the region with a length (along the x -axis) equal to the laser pulse length, and the width (along the y -axis) equal to $l_{ch} \tan \theta_{eff}$. This region is co-moving with the fundamental mode inside the fibre. The amplitude of the n -th harmonic can be estimated as $(a_0 \sin \theta_{eff})^n (\omega_0/\omega_{pe})^{1+2n}$, where $\omega_{pe} = (4\pi n_0 e^2/m_e)^{1/2}$. For $a_0 = 3$, $(\omega_0/\omega_{pe})^2 = 1/8$ and $\theta_{eff} \approx 38^\circ$, we obtain $E^{(n)}/E^{(0)} \approx 0.35(0.23)^n$. This proves an effective mechanism for generation of ultra-short pulses of coherent light with a very short wavelength. The corresponding device can be called “the relativistic whistle”.

-
- [1] N. Bloembergen, Nonlinear Optics (Mass.: Reading, 1965); Y. R. Shen, The Principles of Nonlinear Optics (Wiley, New York, 1984); S. A. Akhmanov, R. V. Khokhlov, Problems of Nonlinear Optics (VINITI, Moscow, 1965); R. W. Boyd, Nonlinear Optics (Academic, San Diego, 1992).
 - [2] J. Zhou, J. Peatross, M. M. Murnane, H. C. Kapteyn, I. P. Christov, Phys. Rev. Lett. **76**, 752 (1996); Y. Kobayashi, T. Sekikawa, Y. Nabekawa, S. Watanabe, Opt. Lett. **23**, 64 (1998); P. Villorresi, et al., Phys. Rev. Lett. **85**, 2494 (2000); C. Altucci, et al., Phys. Rev. A **61**, 021801(R) (1999).
 - [3] S. V. Bulanov, F. Califano, G. I. Dudnikova, et al., in: Reviews of Plasma Physics (Kluwer Academic. Plenum Publishers,

New York, 2001), Ed. by V. D. Shafranov, Vol. 22, p. 227.

- [4] G. Mourou, Z. Chang, A. Maksimchuk, et al., Plasma Phys. Rep. **28**, 12 (2002).
- [5] M. Tatarakis, I. Watts, E. N. Beg, et al., Nature **415**, 280 (2002); M. Tatarakis, A. Gopal, I. Watts, et al., Phys. Plasmas **9**, 2244 (2002).
- [6] S. V. Bulanov, N. M. Naumova, F. Pegoraro, Phys. Plasmas **1**, 745 (1994).
- [7] P. Gibbon, Phys. Rev. Lett. **76**, 50 (1996).
- [8] R. Lichters, J. Meyer-ter-Vehn, A. M. Pukhov, Phys. Plasmas **3**, 3425 (1996).
- [9] A. V. Vshivkov, N. M. Naumova, F. Pegoraro, S. V. Bulanov, Phys. Plasmas **5**, 2752 (1998).
- [10] A. Macchi, F. Cornolti, F. Pegoraro, T. V. Liseikina, H. Ruhl, V. A. Vshivkov, Phys. Rev. Lett. **87**, 205004 (2001); A. Macchi, F. Cornolti, F. Pegoraro, Phys. Plasmas **9**, 1704 (2002).
- [11] M. Zepf, G. D. Tsakiris, G. Pretzler, I. Watts, D. M. Chambers, P. A. Norreys, U. Andiel, A. E. Dangor, K. Eidmann, C. Gahn, A. Machacek, J. S. Wark, K. Witte, Phys. Rev. E **58**, 5253 (1998).
- [12] A. Tarasevitch, A. Orisch, D. von der Linde, P. Balcou, G. Rey, J.-P. Chambaret, U. Teubner, D. Klöpfel, W. Theobald, Phys. Rev. E **62**, 023816-1(2000); I. Watts, et al., Phys. Rev. Lett. **88**, 155001 (2002).
- [13] S. V. Bulanov, F. F. Kamenets, A. M. Pukhov, F. Pegoraro, Physics Letters A **195**, 84 (1994).
- [14] A. Bourdier, Phys. Fluids **26**, 1804 (1983).
- [15] T. Zh. Esirkepov, Comput. Phys. Comm. **135**, 144 (2001).